

# Thymoma with Paraneoplastic Syndromes, Good's Syndrome, and Pure Red Cell Aplasia

Jacolien van der Marel, MD, Paul V.M. Pahlplatz, MD, Willem-Hans Steup, MD, PhD,  
and Eduard R. Hendriks, MD

(*J Thorac Oncol.* 2007;2: 325–326)

## Case Report

A 56-year-old man was examined for a 1.5-year history of recurrent airway infections and sinusitis. He had also been diagnosed with pure red cell aplasia (PRCA), hemoglobin concentration 4.7 mmol/l, for which he was treated with corticosteroids (prednisone 1 mg/kg/day) for 1 year and blood transfusions.<sup>1</sup> This treatment stabilized his hemoglobin concentration at 9.5 mmol/l. The recurrent airway infections were first thought to be treatment-related complications by corticosteroids and were treated with antibiotics. Sputum cultures grew *Haemophilus influenzae*. A computed tomographic scan to exclude bronchiectasis revealed a non-invasive thymoma measuring 11 × 5 cm (Figure 1).

Because the sinopulmonary infections did not diminish, additional tests were performed to exclude an immunodeficiency that can occur in the presence of a thymoma. The immunoglobulin levels were very low (IgG 3.2 g/l; IgA and IgM non-detectable), and the patient was diagnosed with Good's syndrome.<sup>2</sup> Treatment with intravenous gamma globulin was started every 6 weeks, which increased the IgG level to 10.2 g/l and led to less frequent pulmonary infections. The noninvasive and encapsulated thymoma was radically re-

sected by median sternotomy<sup>3</sup> (Figure 2). Postoperative recovery was uneventful. Histology confirmed the diagnosis of thymoma, type A/B. As expected, the hypogammaglobulinaemia did not diminish. Therefore, intravenous gamma globulin therapy was continued every 6 weeks. The hematoipoiesis, however, completely recovered after thymectomy.

Thymomas are associated with different paraneoplastic syndromes; myasthenia gravis, Lambert-Eaton myasthenic syndrome, pemphigus, subacute sensory neuropathy, pure red cell aplasia, and immunodeficiency. Most of these are autoimmune or endocrine-related. The most clinically important are myasthenia gravis, PRCA, and hypogammaglobulinaemia. In the presence of a thymoma, PRCA occurs in 5% to 15% and hypogammaglobulinaemia in less than 5% of the cases. Treatment of PRCA consists of thymectomy. However, this may result in normalization of the bone marrow in only 30% to 40%. A recent case series of 12 patients with thymoma and PRCA reported no remission of anemia after thymectomy.<sup>4</sup> In contrast to other paraneoplastic disorders, hypogammaglobulinaemia is not influenced by thymectomy.

## REFERENCES

1. Murakawa T, Nakajima J, Sato H, et al. Thymoma associated with pure red-cell aplasia: clinical features and prognosis. *Asian Cardiovasc Thorac Ann* 2002;10:150–154.
2. Miyakis S, Pefanis A, Passam FH, et al. Thymoma with immunodeficiency (Good's syndrome): review of the literature apropos three cases. *Scand J Infect Dis* 2006;38:314–319.
3. Wright CD, Kessler KA. Surgical treatment of thymic tumors. *Semin Thorac Cardiovasc Surg* 2005;17:20–26.
4. Thompson CA, Steensma DP. Pure red cell aplasia associated with thymoma: clinical insights from a 50-year single-institution experience. *Br J Haematol* 2006;135:405–407.


HagaHospital, Department of Surgery, The Hague, The Netherlands.

Disclosure: The authors declare no conflict of interest.


Address for correspondence: Eduard R. Hendriks, MD, HagaHospital, Department of Surgery, Leyweg 275, 2545CH The Hague, The Netherlands.  
E-mail: e.hendriks@hagaziekenhuis.nl

Copyright © 2007 by the International Association for the Study of Lung Cancer

ISSN: 1556-0864/07/0204-0325



**FIGURE 1.** Chest computed tomographic scan showing a large mediastinal mass (thymoma). *A*, Axial view. *B*, Coronal view.



**FIGURE 2.** Radical resected thymoma. Type A/B, mixed thymoma. Size 11 × 5 cm.